

ON CHLODOWSKY VARIANT OF (p, q) KANTOROVICH-STANCU-SCHURER OPERATORS

VISHNU NARAYAN MISHRA^{1,2,*} AND SHIKHA PANDEY¹

ABSTRACT. In the present paper, we introduce the Chlodowsky variant of (p, q) Kantorovich-Stancu-Schurer operators on the unbounded domain which is a generalization of (p, q) Bernstein-Stancu-Kantorovich operators. We have also derived its Korovkin type approximation properties and rate of convergence.

1. INTRODUCTION AND PRELIMINARIES

Approximation theory has an important role in mathematical research because of its great potential for applications. Korovkin gave his famous approximation theorem in 1950, since then the study of the linear methods of approximation given by sequences of positive and linear operators became a deep-rooted part of approximation theory. Considering it, various operators as Bernstein, Szász, Baskakov etc. and their generalizations are being studied. In recent years, many results about the generalization of positive linear operators have been obtained by several mathematicians ([6], [8]-[12], [21]). In last two decades, the applications of q -calculus has played an important role in the area of approximation theory, number theory and theoretical physics. In 1987, Lupaş and in 1997, Phillips introduced a sequence of Bernstein polynomials based on q -integers and investigated its approximation properties. Several researchers obtained various other generalizations of operators based on q -calculus(See [3], [17]-[19]).

Recently, Mursaleen et al. applied (p, q) -calculus in approximation theory and introduced first (p, q) -analogue of Bernstein operators. They investigated uniform convergence of the operators and order of convergence, obtained Voronovskaja theorem as well. Also, (p, q) -analogue of Bernstein operators, Bernstein-Stancu operators and Bernstein-Schurer operators, Kantorovich Bernstein Schurer, and Bleimann-Butzer-Hahn operators were introduced in ([13]-[16]), respectively. Further, Acar [1] have studied recently, (p, q) -Generalization of SzászMiraky operators.

In the present paper, we introduce the Chlodowsky variant of (p, q) Kantorovich-Stancu-Schurer operators on the unbounded domain. We begin by recalling certain notations of (p, q) -calculus.

For $0 < q < p \leq 1$, the (p, q) integer $[n]_{p,q}$ is defined by

$$[n]_{p,q} := \frac{p^n - q^n}{p - q}.$$

(p, q) factorial is expressed as

$$[n]_{p,q}! = [n]_{p,q}[n-1]_{p,q}[n-2]_{p,q} \dots 1.$$

(p, q) binomial coefficient is expressed as

$$\left[\begin{array}{c} n \\ k \end{array} \right]_{p,q} := \frac{[n]_{p,q}!}{[k]_{p,q}![n-k]_{p,q}!}.$$

2010 Mathematics Subject Classification. 41A25, 41A36, 41A10, 41A30.

Key words and phrases. (p, q) -integers; (p, q) Bernstein operators; Chlodowsky polynomials; (p, q) Bernstein-Kantorovich operators; modulus of continuity; linear positive operator; Korovkin type approximation theorem; rate of convergence.

©2016 Authors retain the copyrights of their papers, and all open access articles are distributed under the terms of the Creative Commons Attribution License.

(p, q) binomial expansion as

$$(ax + by)_{p,q}^n := \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} a^{n-k} b^k x^{n-k} y^k.$$

$$(x + y)_{p,q}^n := (x + y)(px + qy)(p^2x + q^2y) \dots (p^{n-1}x + q^{n-1}y).$$

The definite integrals of the function f are given by

$$\int_0^a f(x) d_{p,q} x = (q-p)a \sum_{k=0}^{\infty} \frac{p^k}{q^{k+1}} f\left(\frac{p^k}{q^{k+1}} a\right), \quad \left|\frac{p}{q}\right| < 1,$$

and

$$\int_0^a f(x) d_{p,q} x = (p-q)a \sum_{k=0}^{\infty} \frac{q^k}{p^{k+1}} f\left(\frac{q^k}{p^{k+1}} a\right), \quad \left|\frac{p}{q}\right| > 1.$$

Further (p, q) analysis can be found in [2].

In 1932, Chlodowsky [7] presented a generalization of Bernstein polynomials on an unbounded set, known as Bernstein-Chlodowsky polynomials given by,

$$(1.1) \quad B_n(f, x) = \sum_{k=0}^n f\left(\frac{k}{n} b_n\right) \binom{n}{k} \left(\frac{x}{b_n}\right)^k \left(1 - \frac{x}{b_n}\right)^{n-k}, \quad 0 \leq x \leq b_n,$$

where b_n is an increasing sequence of positive terms with the properties $b_n \rightarrow \infty$ and $\frac{b_n}{n} \rightarrow 0$ as $n \rightarrow \infty$.

In 2015, Vedi and Özarslan [20] investigated Chlodowsky-type q -Bernstein-Stancu-Kantorovich operators, and Wafi and Rao investigated (p, q) form of Kantorovich type Bernstein-Stancu-Schurer operator. Mursaleen and Khan [15] defined the Kantorovich type (p, q) -Bernstein-Schurer Operators, given by

$$T_{n,m}(f; x, p, q) = \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} x^k \prod_{s=0}^{n+m-k-1} (p^s - q^s x) \int_0^1 f\left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t}{[n+1]_{p,q}}\right) d_{p,q} t$$

$$k = 0, 1, 2, \dots, n = 1, 2, 3, \dots$$

where

Lemma 1. (See [15]) For the Operators $T_{n,m}^{(\alpha, \beta)}$, we have

$$T_{n,m}(1; x, p, q) = 1,$$

$$T_{n,m}(t; x, p, q) = \frac{(px + 1 - x)_{p,q}^{n+m}}{[2]_{p,q}[n+1]_{p,q}} + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}[n+1]_{p,q}} x,$$

$$T_{n,m}(t^2; x, p, q) = \frac{(p^2x + 1 - x)_{p,q}^{n+m}}{[3]_{p,q}[n+1]_{p,q}^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{[n+1]_{p,q}^2} (px + 1 - x)_{p,q}^{n+m-1} x$$

$$+ \left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{[n+1]_{p,q}^2} x^2.$$

2. CONSTRUCTION OF THE OPERATORS

We construct the Chlodowsky variant of (p, q) Kontorovich-Stancu-Schurer operators as

$$(2.1) \quad K_{n,m}^{(\alpha, \beta)}(f; x, p, q) = \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n}\right)^k \int_0^1 f\left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t + \alpha b_n}{[n+1]_{p,q} + \beta}\right) d_{p,q} t,$$

where $n \in \mathbb{N}$, $m, \alpha, \beta \in \mathbb{N}_0$ with $0 \leq \alpha \leq \beta$, $0 \leq x \leq b_n$, $0 < q < p \leq 1$.

Obviously, $K_{n,m}^{(\alpha, \beta)}$ is a linear and positive operator. To begin with, we obtain the following important lemma.

Lemma 2. Let $K_{n,m}^{(\alpha,\beta)}(f; x, p, q)$ be given by (2.1). The first few moments of the operators are

- (i) $K_{n,m}^{(\alpha,\beta)}(1; x, p, q) = 1,$
- (ii) $K_{n,m}^{(\alpha,\beta)}(t; x, p, q) = \left(\frac{1}{[n+1]_{p,q} + \beta} \right) \left(\alpha b_n + \frac{(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}} x \right),$
- (iii)
$$\begin{aligned} K_{n,m}^{(\alpha,\beta)}(t^2; x, p, q) \\ = \left(\frac{1}{[n+1]_{p,q} + \beta} \right)^2 \left[\left(\alpha^2 + \frac{2\alpha}{[2]_{p,q}} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}} \right) b_n^2 \right. \\ \left. + \left(\frac{2\alpha}{[2]_{p,q}} (p+2q-1) + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right) [n+m]_{p,q} b_n x \right. \\ \left. + \left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} [n+m]_{p,q} [n+m-1]_{p,q} x^2 \right], \end{aligned}$$
- (iv) $K_{n,m}^{(\alpha,\beta)}((t-x); x, p, q) = \frac{[2]_{p,q} \alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} - 1 \right) x$
- (v)
$$\begin{aligned} K_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) \\ = \left[\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q} ([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q} ([n+1]_{p,q} + \beta)^2} \right] b_n^2 \\ + \left[\frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right. \\ \left. - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} \right] b_n x \\ + \left[\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q} [n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} + 1 \right] x^2. \end{aligned}$$

Proof. From operator 2.1,

$$\begin{aligned}
K_{n,m}^{(\alpha,\beta)}(t^u; x, p, q) &= \sum_{k=0}^{n+m} \left[\begin{array}{c} n+m \\ k \end{array} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t + \alpha}{[n+1]_{p,q} + \beta} b_n \right)^u d_{p,q} t \\
&= \sum_{k=0}^{n+m} \left[\begin{array}{c} n+m \\ k \end{array} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \frac{[n+1]_{p,q}^u b_n^u}{([n+1]_{p,q} + \beta)^u} \\
&\quad \times \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t + \alpha}{[n+1]_{p,q}} \right)^u d_{p,q} t \\
&= \frac{[n+1]_{p,q}^u b_n^u}{([n+1]_{p,q} + \beta)^u} \sum_{k=0}^{n+m} \left[\begin{array}{c} n+m \\ k \end{array} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \sum_{i=0}^u \binom{u}{i} \left(\frac{\alpha}{[n+1]_{p,q}} \right)^{u-i} \\
&\quad \times \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t}{[n+1]_{p,q}} \right)^u d_{p,q} t \\
&= \frac{[n+1]_{p,q}^u b_n^u}{([n+1]_{p,q} + \beta)^u} \sum_{i=0}^u \binom{u}{i} \left(\frac{\alpha}{[n+1]_{p,q}} \right)^{u-i} \sum_{k=0}^{n+m} \left[\begin{array}{c} n+m \\ k \end{array} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \\
&\quad \times \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q} t}{[n+1]_{p,q}} \right)^i d_{p,q} t \\
(2.2) \quad K_{n,m}^{(\alpha,\beta)}(t^u; x, p, q) &= \frac{[n+1]_{p,q}^u b_n^u}{([n+1]_{p,q} + \beta)^u} \sum_{i=0}^u \binom{u}{i} \left(\frac{\alpha}{[n+1]_{p,q}} \right)^{u-i} T_{n,m}(t^i; \frac{x}{b_n}, p, q).
\end{aligned}$$

Thus for $u=0,1,2$ we get

$$K_{n,m}^{(\alpha,\beta)}(1; x, p, q) = T_{n,m}(1; \frac{x}{b_n}, p, q),$$

$$\begin{aligned}
K_{n,m}^{(\alpha,\beta)}(t; x, p, q) &= \frac{[n+1]_{p,q} b_n}{[n+1]_{p,q} + \beta} \sum_{i=0}^1 \binom{1}{i} \left(\frac{\alpha}{[n+1]_{p,q}} \right)^{1-i} T_{n,m}(t^i; \frac{x}{b_n}, p, q) \\
&= \left(\frac{[n+1]_{p,q}}{[n+1]_{p,q} + \beta} \right) b_n \left\{ \frac{\alpha}{[n+1]_{p,q}} + T_{n,m}(t; \frac{x}{b_n}, p, q) \right\}, \\
K_{n,m}^{(\alpha,\beta)}(t^2; x, p, q) &= \frac{[n+1]_{p,q}^2 b_n^2}{([n+1]_{p,q} + \beta)^2} \sum_{i=0}^2 \binom{2}{i} \left(\frac{\alpha}{[n+1]_{p,q}} \right)^{2-i} T_{n,m}(t^i; \frac{x}{b_n}, p, q) \\
&= \frac{[n+1]_{p,q}^2 b_n^2}{([n+1]_{p,q} + \beta)^2} \left[\left(\frac{\alpha}{[n+1]_{p,q}} \right)^2 + 2 \left(\frac{\alpha}{[n+1]_{p,q}} \right) T_{n,m}(t; \frac{x}{b_n}, p, q) + T_{n,m}(t^2; \frac{x}{b_n}, p, q) \right].
\end{aligned}$$

Using Lemma 1 and in view of the above relations we get the statements (i), (ii) and (iii).

Using linear property of operators, we have

$$\begin{aligned}
K_{n,m}^{(\alpha,\beta)}((t-x); x, p, q) &= K_{n,m}^{(\alpha,\beta)}(t; x, p, q) - x K_{n,m}^{(\alpha,\beta)}(1; x, p, q) \\
&= \frac{[2]_{p,q} \alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} - 1 \right) x.
\end{aligned}$$

Hence, we get (iv).

Similar calculations give

$$K_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) = K_{n,m}^{(\alpha,\beta)}(t^2; x, p, q) - 2x K_{n,m}^{(\alpha,\beta)}(t; x, p, q) + x^2 K_{n,m}^{(\alpha,\beta)}(1; x, p, q).$$

Then we obtain,

$$\begin{aligned}
&K_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) \\
&= \left[\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q} ([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q} ([n+1]_{p,q} + \beta)^2} b_n^2 \right. \\
&\quad + \left[\frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right. \\
&\quad \left. - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} \right] b_n x \\
&\quad + \left[\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q} ([n+1]_{p,q} + \beta)} + 1 \right] x^2.
\end{aligned}$$

This proves (v). \square

3. KOROVKIN-TYPE APPROXIMATION THEOREM

Assume C_ρ is the space of all continuous functions f such that

$$|f(x)| \leq M\rho(x), \quad -\infty < x < \infty.$$

Then C_ρ is a Banach space with the norm

$$\|f\|_\rho = \sup_{-\infty < x < \infty} \frac{|f(x)|}{\rho(x)}.$$

The subsequent results are used for proving Korovkin approximation theorem on unbounded sets.

Theorem 1. (See [5]) *There exists a sequence of positive linear operators \mathbb{U}_n , acting from C_ρ to C_ρ , satisfying the conditions*

- (1) $\lim_{n \rightarrow \infty} \|\mathbb{U}_n(1; x) - 1\|_\rho = 0$,
- (2) $\lim_{n \rightarrow \infty} \|\mathbb{U}_n(\phi; x) - \phi\|_\rho = 0$,
- (3) $\lim_{n \rightarrow \infty} \|\mathbb{U}_n(\phi^2; x) - \phi^2\|_\rho = 0$,

where $\phi(x)$ is a continuous and increasing function on $(-\infty, \infty)$ such that $\lim_{x \rightarrow \pm\infty} \phi(x) = \pm\infty$ and $\rho(x) = 1 + \phi^2$, and there exists a function $f^* \in C_\rho$ for which $\lim_{n \rightarrow \infty} \|\mathbb{U}_n f^* - f^*\|_\rho > 0$.

Theorem 2. (See [5]) Conditions (1),(2),(3) of above theorem implies that

$$\lim_{n \rightarrow \infty} \|\mathbb{U}_n f - f\|_\rho = 0$$

for any function f belonging to the subset $C_\rho^0 := \{f \in C_\rho : \lim_{|x| \rightarrow \infty} \frac{|f(x)|}{\rho(x)} \text{ is finite}\}$.

Consider the weight function $\rho(x) = 1 + x^2$ and operators:

$$\mathbb{U}_{n,m}^{\alpha,\beta}(f; x, p, q) = \begin{cases} K_{n,m}^{\alpha,\beta}(f; x, p, q) & \text{if } x \in [0, b_n], \\ f(x) & \text{if } x \in [0, \infty)/[0, b_n]. \end{cases}$$

Thus for $f \in C_{1+x^2}$, we have

$$\begin{aligned} \|\mathbb{U}_{n,m}^{\alpha,\beta}(f; \cdot, p, q)\| &\leq \sup_{x \in [0, b_n]} \frac{|\mathbb{U}_{n,m}^{\alpha,\beta}(f; x, p, q)|}{1 + x^2} + \sup_{b_n < x < \infty} \frac{|f(x)|}{1 + x^2} \\ &\leq \|f\|_{1+x^2} \left[\sup_{x \in [0, \infty)} \frac{|\mathbb{U}_{n,m}^{\alpha,\beta}(1+t^2; x, p, q)|}{1 + x^2} + 1 \right]. \end{aligned}$$

Now, using Lemma 2, we will obtain,

$$\|\mathbb{U}_{n,m}^{\alpha,\beta}(f; \cdot, p, q)\|_{1+x^2} \leq M \|f\|_{1+x^2}$$

if $p := (p)_n, q := (q)_n$ with $0 < q_n < p_n \leq 1, \lim_{n \rightarrow \infty} p_n = 1, \lim_{n \rightarrow \infty} q_n = 1, \lim_{n \rightarrow \infty} p_n^n = \lim_{n \rightarrow \infty} q_n^n = N, N < \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n}{[n]} = 0$.

Theorem 3. For all $f \in C_{1+x^2}^0$, we have

$$\lim_{n \rightarrow \infty} \|\mathbb{U}_{n,m}^{\alpha,\beta}(f; \cdot, p_n, q_n) - f(\cdot)\|_{1+x^2} = 0$$

provided that $p := (p)_n, q := (q)_n$ with $0 < q_n < p_n \leq 1, \lim_{n \rightarrow \infty} p_n = 1, \lim_{n \rightarrow \infty} q_n = 1, \lim_{n \rightarrow \infty} p_n^n = \lim_{n \rightarrow \infty} q_n^n = N, N < \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n}{[n]} = 0$.

Proof. Using the results of Theorem 1 and Lemma 2(i),(ii) and (iii), we will achieve the following assessments, respectively:

$$\begin{aligned} \sup_{x \in [0, \infty)} \frac{|\mathbb{U}_{n,m}^{\alpha,\beta}(1; x, p_n, q_n) - 1|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|K_{n,m}^{\alpha,\beta}(1; x, p_n, q_n) - 1|}{1 + x^2} = 0 \\ \sup_{x \in [0, \infty)} \frac{|\mathbb{U}_{n,m}^{\alpha,\beta}(t; x, p_n, q_n) - t|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|K_{n,m}^{\alpha,\beta}(t; x, p_n, q_n) - x|}{1 + x^2} \\ &\leq \sup_{0 \leq x \leq b_n} \frac{\frac{[2]_{p,q} \alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right) x}{1 + x^2} \\ &\leq \frac{[2]_{p,q} \alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \left| \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right| \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \sup_{x \in [0, \infty)} \frac{|\mathbb{U}_{n,m}^{\alpha,\beta}(t^2; x, p_n, q_n) - t^2|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|K_{n,m}^{\alpha,\beta}(t^2; x, p_n, q_n) - x^2|}{1 + x^2} \\ &\leq \sup_{0 \leq x \leq b_n} \frac{1}{1 + x^2} \left(\left[\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}([n+1]_{p,q} + \beta)^2} \right] b_n^2 \right. \\ &\quad \left. + \left[\frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right] \right) \end{aligned}$$

$$\begin{aligned}
& - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} \Big] b_n x \\
& + \left[\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} + 1 \right] x^2 \\
& \leq \left(\left[\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}([n+1]_{p,q} + \beta)^2} \right] b_n^2 \right. \\
& + \left| \frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right. \\
& - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} \Big| b_n \\
& \left. + \left| \left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} + 1 \right| \right) \rightarrow 0
\end{aligned}$$

whenever $n \rightarrow \infty$, because we have $\lim_{n \rightarrow \infty} p_n = \lim_{n \rightarrow \infty} q_n = 1$ and $\frac{b_n}{[n]_{p,q}} = 0$ as $n \rightarrow \infty$. \square

Theorem 4. Assuming C as a positive and real number independent of n and f as a continuous function which vanishes on $[C, \infty)$. Let $p := (p_n), q := (q_n)$ with $0 < q_n < p_n \leq 1, \lim_{n \rightarrow \infty} p_n = \lim_{n \rightarrow \infty} q_n = 1, \lim_{n \rightarrow \infty} p_n^n = \lim_{n \rightarrow \infty} q_n^n = N < \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n^2}{[n]_{p,q}} = 0$. Then we have

$$\lim_{n \rightarrow \infty} \sup_{0 \leq x \leq b_n} |K_{n,m}^{\alpha,\beta}(f; x, p_n, q_n) - f(x)| = 0.$$

Proof. From the hypothesis on f , it is bounded i.e. $|f(x)| \leq M$ ($M > 0$). For any $\epsilon > 0$, we have

$$\left| f \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right) - f(x) \right| < \epsilon + \frac{2M}{\delta^2} \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2,$$

where $x \in [0, b_n]$ and $\delta = \delta(\epsilon)$ are independent of n . Now since we know,

$$K_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p_n, q_n) = \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right)^2 dt.$$

we can conclude by Theorem 3,

$$\begin{aligned}
& \sup_{0 \leq x \leq b_n} |K_{n,m}^{\alpha,\beta}(f; x, p_n, q_n) - f(x)| \\
& \leq \epsilon + \frac{2M}{\delta^2} b_n^2 \left(\left[\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}([n+1]_{p,q} + \beta)^2} \right] \right. \\
& + \left| \frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right. \\
& - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} \Big| \\
& \left. + \left| \left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} + 1 \right| \right).
\end{aligned}$$

Since $\frac{b_n^2}{[n]_{p,q}} = 0$ as $n \rightarrow \infty$, we have the desired result. \square

4. RATE OF CONVERGENCE

We will find the rate of convergence in terms of the Lipschitz class $Lip_M(\gamma)$, for $0 < \gamma \leq 1$. Assume that $C_B[0, \infty)$ denote the space of bounded continuous functions on $[0, \infty)$. A function $f \in C_B[0, \infty)$ belongs to $Lip_M(\gamma)$ if

$$|f(t) - f(x)| \leq M|t - x|^\gamma, \quad t, x \in [0, \infty)$$

is satisfied.

Theorem 5. Let $f \in Lip_M(\gamma)$, then

$$K_{n,m}^{\alpha,\beta}(f; x, p, q) \leq M(\mu_{n,p,q}(x))^{\gamma/2},$$

where $\mu_{n,p,q}(x) = K_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q)$.

Proof. Since $f \in Lip_M(\gamma)$, and the operator $K_{n,m}^{\alpha,\beta}(f; x, p, q)$ is linear and monotone,

$$\begin{aligned} & |K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\ &= \left| \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \right. \\ &\quad \times \int_0^1 \left(f \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right) - f(x) \right) d_{p,q} t \Big| \\ &\leq \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \\ &\quad \times \int_0^1 \left| f \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right) - f(x) \right| d_{p,q} t \\ &\leq M \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \\ &\quad \times \int_0^1 \left| \frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right|^\gamma d_{p,q} t. \end{aligned}$$

Applying Hölder's inequality with the values $p = \frac{2}{\gamma}$ and $q = \frac{2}{2-\gamma}$, we get following inequality,

$$\begin{aligned} & \int_0^1 \left| \frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right|^\gamma d_{p,q} t \\ &\leq \left\{ \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t \right\}^{\frac{\gamma}{2}} \left\{ \int_0^1 d_{p,q} t \right\}^{\frac{2-\gamma}{2}} \\ &= \left\{ \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t \right\}^{\frac{\gamma}{2}} \end{aligned}$$

Using this, we get

$$\begin{aligned} & |K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\ &\leq M \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k \\ &\quad \times \left\{ \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t \right\}^{\frac{\gamma}{2}} \\ &= M \sum_{k=0}^{n+m} \left\{ \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t \right\}^{\frac{\gamma}{2}} w_{n,k}(p, q; x), \end{aligned}$$

where $w_{n,k}(p, q; x) = \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{x}{b_n} \right)^k$. Again using Hölder's inequality with $p = \frac{2}{\gamma}$ and $q = \frac{2}{2-\gamma}$, we have

$$\begin{aligned} & |K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\ &\leq M \left\{ \sum_{k=0}^{n+m} \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t w_{n,k}(p, q; x) \right\}^{\frac{\gamma}{2}} \left\{ \sum_{k=0}^{n+m} w_{n,k}(p, q; x) \right\}^{\frac{2-\gamma}{2}} \end{aligned}$$

$$\begin{aligned}
&= M \left\{ \sum_{k=0}^{n+m} w_{n,k}(p, q; x) \int_0^1 \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n - x \right)^2 d_{p,q} t \right\}^{\frac{\gamma}{2}} \\
&= M(\mu_{n,p,q}(x))^{\gamma/2},
\end{aligned}$$

where $(\mu_{n,p,q}(x))^{\gamma/2} = K_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q)$. \square

In order to obtain rate of convergence in terms of modulus of continuity $\omega(f; \delta)$, we assume that for any uniformly continuous $f \in C_B[0, \infty)$ and $x \geq 0$, modulus of continuity of f is given by

$$(4.1) \quad \omega(f; \delta) = \max_{\substack{|t-x| \leq \delta \\ t, x \in [0, \infty)}} |f(t) - f(x)|.$$

Thus it implies for any $\delta > 0$

$$(4.2) \quad |f(x) - f(y)| \leq \omega(f; \delta) \left(\frac{|x-y|}{\delta} + 1 \right),$$

is satisfied.

Theorem 6. *If $f \in C_B[0, \infty)$, we have*

$$|K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq 2\omega(f; \sqrt{\mu_{n,p,q}(x)}),$$

where $\omega(f; \cdot)$ is modulus of continuity of f and $\lambda_{n,p,q}(x)$ be the same as in Theorem 5.

Proof. Using triangular inequality, we get

$$\begin{aligned}
&|K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\
&= \left| \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(f \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right) - f(x) \right) \right| \\
&\leq \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| f \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha}{[n+1]_{p,q} + \beta} b_n \right) - f(x) \right|,
\end{aligned}$$

Now using (4.2) and Hölder's inequality, we get

$$\begin{aligned}
&|K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\
&= \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{|(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha b_n - x|}{\delta} + 1 \right) \omega(f; \delta) \\
&\leq \omega(f; \delta) \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \\
&\quad + \frac{\omega(f; \delta)}{\delta} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha b_n - x}{[n+1]_{p,q} + \beta} \right| \\
&= \omega(f; \delta) + \frac{\omega(f; \delta)}{\delta} \left\{ \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right. \\
&\quad \times \left. \left(\frac{(1-t)[k]_{p,q} + [k+1]_{p,q}t + \alpha b_n - x}{[n+1]_{p,q} + \beta} \right)^2 \right\}^{\frac{1}{2}} \\
&= \omega(f; \delta) + \frac{\omega(f; \delta)}{\delta} \{K_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q)\}^{1/2}.
\end{aligned}$$

Now choosing $\delta = \mu_{n,p,q}(x)$ as in Theorem 5, we have

$$|K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq 2\omega(f; \sqrt{\mu_{n,p,q}(x)}).$$

\square

Now let us denote by $C_B^2[0, \infty)$ the space of all functions $f \in C_B[0, \infty)$ such that $f', f'' \in C_B[0, \infty)$. Let $\|f\|$ denote the usual supremum norm of f . The classical Peetre's K-functional and the second modulus of smoothness of the function $f \in C_B[0, \infty)$ are defined respectively as

$$K(f, \delta) := \inf_{g \in C_B^2[0, \infty)} [\|f - g\| + \delta \|g''\|]$$

and

$$\omega_2(f, \delta) = \sup_{\substack{0 < h < \delta, \\ x, x+h \in I}} |f(x+2h) - 2f(x+h) + f(x)|,$$

where $\delta > 0$. It is known that [see [4], p. 177] there exists a constant $A > 0$ such that

$$(4.3) \quad K(f, \delta) \leq A\omega_2(f, \delta).$$

Theorem 7. *Let $x \in [0, b_n]$ and $f \in C_B[0, \infty)$. Then, for fixed $p \in \mathbb{N}_0$, we have*

$$|K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq C\omega_2(f, \sqrt{\alpha_{n,p,q}(x)}) + \omega(f, \beta_{n,p,q}(x))$$

for some positive constant C , where

$$(4.4) \quad \begin{aligned} \alpha_{n,p,q}(x) = & \left[\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} + \frac{(p+2q-1)^2}{[2]_{p,q}^2} \right\} \frac{[n+m]_{p,q}^2}{([n+1]_{p,q} + \beta)^2} - 4 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} + 2 \right] x^2 \\ & + \left[\left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} + 2 \frac{(p+2q-1)}{[2]_{p,q}^2} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} \right. \\ & \quad \left. + 4 \frac{\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} - 4 \frac{(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 4 \frac{\alpha}{([n+1]_{p,q} + \beta)} \right] b_n x \\ & + \left[\frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}} + \frac{(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{2n+2m}}{[2]_{p,q}^2} + 4 \frac{\alpha}{[2]_{p,q}} (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m} + 2\alpha^2 \right] \frac{b_n^2}{([n+1]_{p,q} + \beta)^2}, \end{aligned}$$

and

$$(4.5) \quad \beta_{n,p,q}(x) = \left(\frac{[2]_{p,q}\alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right) x \right).$$

Proof. Consider an auxiliary operator $K_{n,m}^*(f; x, p, q) : C_B[0, \infty) \rightarrow C_B[0, \infty)$ by

$$(4.6) \quad K_{n,m}^*(f; x, p, q) := K_{n,m}^{\alpha,\beta}(f; x, p, q) - f \left(\frac{[2]_{p,q}\alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right) + f(x).$$

Then by Lemma 2 we get

$$(4.7) \quad \begin{aligned} K_{n,m}^*(1; x, p, q) &= 1, \\ K_{n,m}^*((t-x); x, p, q) &= 0. \end{aligned}$$

For given $g \in C_B[0, \infty)$, it follows by the Taylor formula that

$$g(y) - g(x) = (y-x)g'(x) + \int_x^y (y-u)g''(u) du.$$

Taking into account 4.6 and using 4.7, we get

$$\begin{aligned} |K_{n,m}^*(g; x, p, q) - g(x)| &= |K_{n,m}^*(g(y) - g(x); x, p, q)| \\ &= \left| g'(x)K_{n,m}^*((y-x); x, p, q) + K_{n,m}^* \left(\int_x^y (y-u)g''(u) du; x, p, q \right) \right| \\ &= \left| K_{n,m}^* \left(\int_x^y (y-u)g''(u) du; x, p, q \right) \right| \end{aligned}$$

Then by 4.6

$$\begin{aligned} |K_{n,m}^*(g; x, p, q) - g(x)| &= \left| K_{n,m}^* \left(\int_x^y (y-u)g''(u) du; x, p, q \right) - \int_x^y \left(\frac{[2]_{p,q}\alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right) du \right| \end{aligned}$$

$$\begin{aligned} & \left| \frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x - u \right) g''(u) du \Big| \\ & \leq \left| K_{n,m}^* \left(\int_x^y (y-u) g''(u) du; x, p, q \right) \right| + \left| \int_x^{\left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right)} (y-u) g''(u) du \right. \\ & \quad \left. \left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x - u \right) g''(u) du \right|. \end{aligned}$$

Since,

$$\left| K_{n,m}^{\alpha,\beta} \left(\int_x^y (y-u) g''(u) du; x, p, q \right) \right| \leq \|g''(x)\| K_{n,m}^{\alpha,\beta}((y-x)^2; x, p, q)$$

and

$$\begin{aligned} & \left| \int_x^{\left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right)} (y-u) g''(u) du \right. \\ & \quad \left. \left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x - u \right) g''(u) du \right| \\ & \leq \|g''\| \left[\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right) x \right]^2, \end{aligned}$$

we get

$$\begin{aligned} |K_{n,m}^*(g; x, p, q) - g(x)| & \leq \|g''\| K_{n,m}^{\alpha,\beta}((y-x)^2; x, p, q) + \|g''\| \left[\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n \right. \\ & \quad \left. + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right) x \right]^2. \end{aligned}$$

Hence Lemma 2 implies that

$$\begin{aligned} (4.8) \quad & |K_{n,m}^*(g; x, p, q) - g(x)| \\ & \leq \|g''\| \left[\left(\frac{\alpha^2}{([n+1]_{p,q} + \beta)^2} + \frac{2\alpha}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} \left(p\frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} + \frac{(p^2\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}([n+1]_{p,q} + \beta)^2} \right) b_n^2 \right. \\ & \quad + \left(\frac{2\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)^2} + \left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q} + \beta)^2} \left(p\frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m-1} \right. \\ & \quad \left. - \frac{2\alpha}{([n+1]_{p,q} + \beta)} - \frac{2(p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} \right) b_n x \\ & \quad + \left(\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} \right\} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n+1]_{p,q} + \beta)^2} - 2 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} + 1 \right) x^2 \\ (4.9) \quad & \quad \left. + \left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} - 1 \right) x \right)^2 \right]. \end{aligned}$$

Since $K_{n,m}^*(f; x, p, q) \leq 3\|f\|$, considering 4.4 and 4.5, for all $f \in C_B[0, \infty)$ and $g \in C_B^2[0, \infty)$, we may write from 4.8 that

$$\begin{aligned} & |K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq |K_{n,m}^*(f-g; x, p, q) - (f-g)(x)| + |K_{n,m}^*(g; x, p, q) - g(x)| \\ & \quad + \left| f \left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right) - f(x) \right| \\ & \leq 4\|f-g\| + \alpha_{n,p,q}(x)\|g'\| + \left| f \left(\frac{[2]_{p,q}\alpha + (p\frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q} + \beta)} b_n + \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q} + \beta)} x \right) - f(x) \right| \\ & \leq 4\|f-g\| + \alpha_{n,p,q}(x)\|g'\| + \omega(f, \beta_{n,p,q}(x)), \end{aligned}$$

which yields that

$$\begin{aligned} |K_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| &\leq 4K(f, \alpha_{n,p,q}(x)) + \omega(f, \text{beta}_{n,p,q}(x)) \\ &\leq C\omega_2(f, \sqrt{\alpha_{n,p,q}(x)}) + \omega(f, \beta_{n,p,q}(x)), \end{aligned}$$

where

$$\begin{aligned} \alpha_{n,p,q}(x) &= \left[\left\{ 1 + \frac{2(q-1)}{[2]_{p,q}} + \frac{(q-1)^2}{[3]_{p,q}} + \frac{(p+2q-1)^2}{[2]_{p,q}^2} \right\} \frac{[n+m]_{p,q}^2}{([n+1]_{p,q}+\beta)^2} - 4 \frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q}+\beta)} + 2 \right] x^2 \\ &+ \left[\left\{ 1 + \frac{2q}{[2]_{p,q}} + \frac{q^2-1}{[3]_{p,q}} + 2 \frac{(p+2q-1)}{[2]_{p,q}^2} \right\} \frac{[n+m]_{p,q}}{([n+1]_{p,q}+\beta)^2} \left(p \frac{x}{b_n} + 1 - \frac{x}{b_n} \right)_{p,q}^{n+m} \right. \\ &+ \left. 4 \frac{\alpha(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q}+\beta)^2} - 4 \frac{(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q}+\beta)} - 4 \frac{\alpha}{([n+1]_{p,q}+\beta)} \right] b_n x \\ &+ \left[\frac{(p^2 \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[3]_{p,q}} + \frac{(p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{2n+2m}}{[2]_{p,q}^2} + 4 \frac{\alpha}{[2]_{p,q}} (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m} + 2\alpha^2 \right] \frac{b_n^2}{([n+1]_{p,q}+\beta)^2}, \end{aligned}$$

and

$$\beta_{n,p,q}(x) = \left(\frac{[2]_{p,q}\alpha + (p \frac{x}{b_n} + 1 - \frac{x}{b_n})_{p,q}^{n+m}}{[2]_{p,q}([n+1]_{p,q}+\beta)} b_n + \left(\frac{(p+2q-1)[n+m]_{p,q}}{[2]_{p,q}([n+1]_{p,q}+\beta)} - 1 \right) x \right).$$

Hence we get the result. \square

REFERENCES

- [1] T. Acar, (p, q) -Generalization of Szász-Mirakyan operators, *Math. Meth. Appl. Sci.*, 2015 (2015), DOI: 10.1002/mma.3721.
- [2] I. M. Burban, A. U. Klimyk, (p, q) -differentiation, (p, q) -integration, and (p, q) -hypergeometric functions related to quantum groups, *Integral Transforms and Special Functions*, 2 (1994), 15-36.
- [3] N. L. Braha, H. M. Srivastava and S. A. Mohiuddine, A Korovkin's type approximation theorem for periodic functions via the statistical summability of the generalized de la Vallée Poussin mean, *Appl. Math. Comput.* 228 (2014), 162-169.
- [4] R.A. Devore, G.G. Lorentz, *Constructive Approximation*, Springer, Berlin, 1993.
- [5] A. D. Gadzhiev, The convergence problem for a sequence of positive linear operators on unbounded sets and theorems analogues to that of P.P. Korovkin, *Dokl. Akad. Nauk SSSR*, 218 (5), 1001-1004. English translation in *Sov. Math. Dokl.*, 15 (1974), 1433-1436.
- [6] A.R. Gairola, Deepmala, L.N. Mishra, Rate of Approximation by Finite Iterates of q -Durrmeyer Operators, *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 2016 (2016), doi: 10.1007/s40010-016-0267-z.
- [7] E. Ibikli, Approximation by Bernstein-Chlodowsky polynomials, *Hacettepe Journal of Mathematics and Statistics*, 32 (2003), 1-5.
- [8] V.N. Mishra, K. Khatri, L.N. Mishra, Deepmala; Inverse result in simultaneous approximation by Baskakov-Durrmeyer-Stancu operators, *Journal of Inequalities and Applications*, 2013 (2013), Article ID 586.
- [9] V.N. Mishra, K. Khatri, L.N. Mishra; On Simultaneous Approximation for Baskakov-Durrmeyer-Stancu type operators, *Journal of Ultra Scientist of Physical Sciences*, 24 (2012), 567-577.
- [10] V.N. Mishra, K. Khatri, L.N. Mishra; Statistical approximation by Kantorovich type Discrete q -Beta operators, *Advances in Difference Equations* 2013 (2013), Article ID 345.
- [11] V.N. Mishra, S. Pandey, On (p, q) Baskakov-Durrmeyer-Stancu Operators, arXiv:1602.06719 [math.CA]
- [12] V.N. Mishra, S. Pandey, (p, q) -Szász-Mirakyan-Baskakov-Stancu type Operators, arXiv:1602.06312 [math.CA].
- [13] M. Mursaleen, K.J. Ansari, A. Khan, On (p, q) -analogue of Bernstein operators, *Appl. Math. Comput.*, 266 (2015), 874-882.
- [14] M. Mursaleen, K.J. Ansari, A. Khan, Some approximation results by (p, q) -analogue of Bernstein-Stancu operators, *Appl. Math. Comput.*, 264 (2015), 392-402.
- [15] M. Mursaleen, F. Khan, Approximation by Kantorovich type (p, q) -Bernstein Schurer operators, arXiv:1506.02492 [math.CA].
- [16] M. Mursaleen, Md. Nasiruzzaman, A. Khan, K.J. Ansari, Some approximation results on Bleimann-Butzer-Hahn operators defined by (p, q) -integers, arXiv:1505.00392, [math.CA].
- [17] M. Mursaleen, A. Khan, H. M. Srivastava and K. S. Nisar, Operators constructed by means of q -Lagrange polynomials and A-statistical approximation, *Appl. Math. Comput.* 219 (2013), 6911-6818.
- [18] H. M. Srivastava, Some generalizations and basic (or q) extensions of the Bernoulli, Euler and Genocchi polynomials, *Appl. Math. Inform. Sci.* 5 (2011), 390-444.
- [19] H. M. Srivastava and J. Choi, *Zeta and q -Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

- [20] T. Vedi and Mehmet Ali Özarslan, Chlodowsky-type q -Bernstein-Stancu-Kantorovich operators, *J. Inequal. Appl.*, 2015 (2015), Article ID 91.
- [21] A. Wafi, N. Rao, Deepmala, Approximation properties by generalized-Baskakov-Kantorovich-Stancu type operators, *Appl. Math. Inf. Sci. Lett.*, 4 (2016), 1-8.

¹DEPARTMENT OF APPLIED MATHEMATICS & HUMANITIES, SARDAR VALLABHBHAI NATIONAL INSTITUTE OF TECHNOLOGY, ICHCHHANATH MAHADEV DUMAS ROAD, SURAT -395 007 (GUJARAT), INDIA

²L. 1627 AWADH PURI COLONY BENIGANJ, PHASE -III, OPP. INDUSTRIAL TRAINING INSTITUTE, AYODHYA MAIN ROAD, FAIZABAD-224 001, (UTTAR PRADESH), INDIA

*CORRESPONDING AUTHOR: VISHNUNARAYANMISHRA@GMAIL.COM